



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

BRIEFER ARTICLES

VARIATION IN HELIANTHUS

In a recent number of this journal, Dr. GEORGE H. SHULL¹ calls attention to the dark disk found in the wild sunflower of the prairie region, but shows that certain material received from that region was heterozygous with a yellow disk recessive. It may be worth while to report that this wild plant does actually produce a yellow-disked variety, which I have observed both in Colorado and New Mexico. The western sunflower, according to RYDBERG, is separable as a distinct species, *Helianthus lenticularis* Dougl. This seems to be going too far, and I think it should be called *H. annuus lenticularis*. The color of the disk is considered a specific character in *Helianthus*; but it certainly varies within specific limits, not only in *H. annuus*, but in *H. petiolaris* also, as I have shown in *Nature*, June 19, 1902, p. 174.—T. D. A. COCKERELL, *Boulder, Colorado*.

ENDOSPERM OF PONTEDERIACEAE

(WITH FOUR FIGURES)

In a recent paper² on the seeds of Pontederiaceae, COKER makes some references to my paper published in 1898,³ to which I wish to make a brief reply. He says: "Oddly enough he completely overlooked the interesting peculiarity in the endosperm of all three genera." The reason for my overlooking such a peculiarity is clear enough, as a very casual reading of the paper will show that I made no attempt to follow endosperm formation. My study was based chiefly on *Eichhornia*, and in that genus, as grown in Washington Park, Chicago, the endosperm nucleus almost never divides, and the contents of the embryo sac disorganize without development of endosperm or embryo.

Some of my slides of *Pontederia*, however, extend beyond the fertilization stage, and since COKER's article appeared I have reexamined these, and am able in three cases to confirm his account of a division of the embryo sac into upper and lower chambers. *Fig. 3* shows the condition of the


¹ Some new cases of Mendelian inheritance. BOT. GAZETTE 45:103-116. 1908.

² COKER, W. C., The development of the seed in the Pontederiaceae. BOT. GAZETTE 44:293-301. *pl.* 23. 1907.

³ SMITH, W. R., A contribution to the life history of the Pontederiaceae. BOT. GAZETTE 25:324-337. *pls.* 19, 20. 1898.

embryo sac just after the division of the endosperm nucleus, and *fig. 4* the lower end of the sac a short time later, with several nuclei in the lower chamber. Of later conditions of the endosperm my slides do not furnish examples.

Again COKER says: "The antipodals never divide, but they are not ephemeral as described by SMITH." There may be some difference of opinion as to what degree of persistence my description implies, but after a reexamination of my slides, and even from COKER's own figures, I am still convinced of its correctness. COKER makes the error of regarding the antipodals as a group of nuclei; he cites the persistence of such nuclei after



the initiation of endosperm formation as a disproof of my statement; but the antipodals are cells, not nuclei. In those embryo sacs in which they are recognized as typically persistent, such as numerous *Ranunculaceae*, *Sparanium*, and others, they become a conspicuous mass of cells; they do not persist as free nuclei. In the case of *Eichhornia* and *Pontederia* they are organized cells at the stage shown in *figs. 1* and *2* herewith, and in *fig. 13* of COKER's plate. These cells disorganize immediately after the fusion of the polar nuclei, sometimes even earlier, a behavior which amply justifies my describing them as ephemeral. In later stages of the embryo sac only the nuclei can be distinguished, and these in a partially disorganized condition, as in *figs. 3* and *4*. Compare also COKER's *figs. 13, 15, 17*, in which the disintegration of the antipodal cells and the absorption of their nuclei may be clearly traced.—R. WILSON SMITH, *McMaster University, Toronto, Canada*.